



Nutrients/food chain model for Lake Zapotlan (Mexico)

MARIO A. ORTIZ-JIMÉNEZ, *Universidad Autónoma de Nayarit, Ciudad de la Cultura, C.P. 63155, Tepic, Nayarit*
E-mail: ojimez@nayar.uan.mx

JOSÉ DE ANDA, *Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco, A.C., Normalistas 800, 44270 Guadalajara, Jalisco, México.* E-mail: janda@ciatej.net.mx

HARVEY SHEAR, *Department of Geography, University of Toronto at Mississauga, 3359 Mississauga Road North, Mississauga, Canada.* E-mail: hshear@utm.utoronto.ca

ABSTRACT

A nutrients/food chain model was developed considering the linkage between water balance and energy balance models for Lake Zapotlan. Lake Zapotlan is located in the southern part of Jalisco State, Mexico, and is classified as a warm, tropical water body, endorheic, very shallow, and highly eutrophic. Kinetic parameters were calibrated using previously reported values in the literature for other lakes. The model simulates the concentration of eight state variables for the lake: algae, herbivore and carnivore zooplankton, nitrates, ammonium, SRP, DOC and POC. A set of eight simultaneous ordinary differential equations (ODE) were generated, assuming an initial set of conditions considering the existing hydrologic and water quality information of the streams flowing into the lake. The ODE system was solved and calibrated by using observed water quality data of the lake. In this way, we have estimated the internal fluxes of nutrients and dissolved oxygen in the lake and their interaction with the food chain.

Keywords: Eutrophication; food chain; flux of nutrients; nutrients load; tropical lakes.

1 Introduction

The Lake Zapotlan Basin is located in the southern part of Jalisco State, Mexico, between 19°34' and 19°53' north latitude and 103°24' and 103°38' west longitude. This basin is the second most populous in Jalisco State, with a population close to one hundred thousand inhabitants [25]. This shallow and endorheic lake shows signs of severe eutrophication, manifested by the presence of large stands of aquatic macrophytes (common cattail [*Typha latifolia*], aquatic hyacinth [*Eichhornia crassipes*]), and periodic blooms of blue-green algae (*Aphanocapsa elachista*, *Chroococcus limneticus*, *Dactylococcopsis acicularis*, and *Microcystis aeruginosa*, between others). The main nutrient sources are from continuous discharges of poorly treated sewage, urban runoff from the cities of Ciudad Guzmán and San Sebastián del Sur, and agricultural and farming runoff around the lake basin [36].

Nutrient/food chain models attempt to characterize the partitioning of matter within the lake on a seasonal basis. These models typically have a number of common characteristics, including transport and kinetic characterization. The variables describing the biological processes include several forms of nutrients and a food chain that relates the different interaction between living microorganisms [46, 50]. These models were primarily developed from the 1970's onwards to address the impact of nutrients on natural waters [4, 9] and include the following

nutrients models: CE-QUAL-W2, a water quality and hydrodynamic model in 2D for water bodies basin systems [12]; PCLAKE, a model that calculates the growth of algae, fish and plants in shallow lakes, within the framework of closed nutrient cycles [28, 26]; a coupled watershed and lake model (AnnAGNPS and BATHYTUB) that simulates the lake water quality conditions of a reservoir considering changes in different watershed land use and management scenarios [27].

In the last two years several water quality models for lakes and reservoirs have also been developed. Some of these are described below:

- (a) A time dependent mathematical model that accounts for the vertical fluxes of nutrients into a lake [23];
- (b) a shallow lake model that takes into account the inherent variability in precipitation and uncertainties in the empirical relationships determining nutrient export from stream catchments [38];
- (c) LAKELOAD, an integrated model that estimates the loadings of nutrients and organic matter from point and non-point sources in a watershed and accounts for the loadings from industrial and municipal wastewaters, atmospheric deposition and runoff from the watershed [40];
- (d) macroscopic food web model that identifies the key and redundant uncertainties associated with the constituent processes of the reservoir ecosystem of Lake Lanier [37];

- (e) a model that applies future scenarios for a lake basin, which are then translated into quantitative changes in the input variables for a GIS system based nutrient transport model [34];
- (f) steady state lake phosphorus mass balance model that is used to predict the equilibrium phosphorus concentration of the lake from natural and anthropogenic, external and internal phosphorus inputs [35];
- (g) MODIFFUS, a nutrient-flow simulation model that estimates nutrient inputs from diffuse sources into surface waters [39];
- (h) a eutrophication model that simulates plankton dynamics in Lake Washington [2];
- (i) a phosphorus management model that aids local officials in decision-making processes pertaining to the management of City Park Lake, a shallow, subtropical, urban hypereutrophic lake [44].

However, due to the shallow and morphometric features of the warm tropical and subtropical Mexican natural lakes, most of the water quality models developed in the literature for temperate and deep lakes and reservoirs fail. In this work, we therefore present the first effort to model a Mexican tropical shallow lake located in a closed basin.

2 Methodology

2.1 Water quantity of Lake Zapotlan

A hydrological balance of Lake Zapotlan was developed considering topo-bathymetric information [5], watershed meteorological data [11], and the initial water level of the lake measured in June of 1982 [5]. A simulation program was also developed to predict the main hydrometric components of the lake for the period between June 1982 and December 2003. This program was developed using a discrete simulation language GPSS (General Purpose Simulation System) [33]. The main output variables

of the hydrologic balance simulation were inflow (runoff, precipitation, groundwater infiltration, and sewage) and outflow of lake (water extraction and evapotranspiration) [36].

2.2 Temperature of the lake

The daily solar radiation and the theoretical hours of monthly insolation in a cloudless sky over lake were calculated. Effective hours of monthly insolation, air vapour pressure, wind velocity, and air temperature over lake were estimated [11]. We obtained a differential equation for temperature that was numerically integrated by the fourth order Runge-Kutta method with a time increment of one day [8].

2.3 State variables and modeling the dynamic of the process

The state variables of the model are divided into three main groups: food chain (algae, herbivorous zooplankton, and carnivorous zooplankton), non-living organic carbon (*POC* and *DOC*), and nutrients ($NH_4 - N$, $NO_3 - N$, and *SRP*) (See Figure 1). The model simulates the competition between phytoplankton and herbivorous and carnivorous zooplankton. The following assumptions were considered in the simulation model:

- (a) algae and detritus are consumed and settle to the bottom of the lake;
- (b) algal growth depends on nutrient concentration, solar radiation, and Lake temperature;
- (c) total zooplankton biomass increases through the consumption of algae and decreases through excretion, respiration and predation by fishes;
- (d) inefficiencies of zooplankton grazing (assumed efficiency of 0.7) provide inputs to the storage of detritus [7];
- (e) the mass balances of the food chain and nutrients considered the Lotka-Volterra and Michaelis-Menten equations, respectively [7],




Figure 1 Nutrient/food-chain model.

(f) because of its shallowness and wind action over the surface, the lake was considered as completely mixed horizontally and vertically. Therefore, the main assumption is that a zero-dimensional model could be a good start to describe the chemical and biological processes occurring in the lake.

We have adopted a mechanistic, deterministic approach for the model, based on the conservation of mass, and on measured input parameters monitored on a single day during each of the dry and rainy seasons, respectively. Therefore, in this first modeling effort, we did not consider the possible variability in these measurements, and how might this variability affect the conclusions drawn. The equation of the simulation model, in explicit form for every state variable v , can be written as follow:

$$\frac{dv}{dt} = \frac{Q_1 c_1}{V} + \frac{Q_2 c_2}{V} + \frac{Q_3 c_3}{V} + \frac{Q_4 c_4}{V} - \frac{Q_5 c}{V} \pm \frac{F_v A}{V \times 1000} + S_v \quad (1)$$

The ratios of chlorophyll-*a*/carbon and nitrogen/carbon were calculated using the Laws-Chalup model [29] and the solar extinction coefficient was calculated using the Riley equation [43]. Ranges of kinetic parameter values reported in the literature [1, 3, 7, 15, 19, 24, 46, 47, 49] were used to simulate the interaction of nutrients/food chain in Lake Zapotlan. The process velocities $k(T)$ were adjusted by temperature with the theta model [7], shown below,

$$k(T) = k_{20} \theta^{T-20} \quad (2)$$

3 Estimation of the state variables

3.1 Inorganic nutrients point sources

Table 1 shows the *TP* and $NO_3 - N$ concentrations observed in the sewage discharges to the lake during the dry season (May 8, 2003). In this kind of lake, when measured *TP* concentrations are higher than 2 mg/L, 80% or more of *TP* is in the form of *SRP* [13, 17]. From the hydraulic balance of the water body [36], it was estimated that direct sewage amounted to about 10,300 m³/d discharged into the lake (Q'_1), and consequently the mass loads of $NO_3 - N$ and *SRP* were, respectively:

$$W'_{n_{i,1}} = Q'_1 n'_{i,1} = 2.88 \times 10^{11} \frac{\mu\text{gN}}{d} \quad (3)$$

$$W'_{p_{s,1}} = Q'_1 p'_{s,1} = 3.87 \times 10^{11} \frac{\mu\text{gP}}{d} \quad (4)$$

Table 1 Nutrient concentrations from direct sewage discharge during the dry season, in mg/L.

Sewage discharges	Observed $NO_3 - N$ ($n'_{i,1}$)	Observed <i>TP</i>	Estimated <i>SRP</i> ($p'_{s,1}$)
Ciudad Guzmán	29.61	47.91	38.33
San Sebastián del Sur	24.11	46.16	36.93
Weighted mean	28.00	47.00	37.60

If the mass load velocities remain constant, $NO_3 - N$ and *SRP* concentrations in the sewage discharges can be written as follows, respectively:

$$n_{i,1} = \frac{2.88 \times 10^{11}}{Q_1 \times 1,000} \quad (5)$$

$$p_{s,1} = \frac{3.87 \times 10^{11}}{Q_1 \times 1,000} \quad (6)$$

3.2 DOC point sources

Table 2 shows the estimated *DOC* in the sewage discharges during the dry season starting from the observed BOD_5 [20]:

$$DOC = \frac{BOD_5 + 5.72}{2.06} \quad (7)$$

3.3 Diffuse sources of nutrients

Table 3 shows the diffuse loads of *SRP* and $NO_3 - N$ from the basin, using the export coefficients proposed by Ryding and Rast [45]. The export coefficients for *SRP* are usually 40 to 50% of that for *TP* [6]. For the purpose of this calculation, we propose that 50% of the *TP* load corresponds to *SRP*. Table 4 shows the diffuse sources of $NO_3 - N$ using the export coefficient for tropical watersheds [31]. Generalized export coefficients are used in this paper due to lack of monitoring information related to loadings from individual land uses in the area of study [32] and consequently we did not consider the likely variability in these coefficients, and how might this variability affect the diffuse loads of nutrients from the basin.

Most of the diffuse loads of *SRP* and $NO_3 - N$ reach the lake through surface runoff [42], and their corresponding concentrations are expressed respectively as:

$$p_{s,2} = \frac{\bar{W}'_{ps}}{\bar{Q}_2} = \frac{43.94 \times 10^{12} \mu\text{gP}}{18.91 \times 10^9 \text{ L}} = 2,323.67 \frac{\mu\text{gP}}{\text{L}} \quad (8)$$

$$n_{i,2} = \frac{\bar{W}'_{ni}}{\bar{Q}_2} = \frac{103.28 \times 10^{12} \mu\text{gN}}{18.91 \times 10^9 \text{ L}} = 5,461.77 \frac{\mu\text{gN}}{\text{L}} \quad (9)$$

3.4 Monitoring of water quality

Table 5 shows the *TP* mean concentration of nine monitoring stations in the lake during the dry and rainy season in 2003. Because we measured only *TP*, it was assumed that 60% of the observed *TP* is in the form of *SRP*, as shown by de Anda *et al.* [14] in Lake Chapala, which has similar water quality features and is located very close to Lake Zapotlan.

Table 2 BOD_5 and *DOC* concentrations from the sewage (mg/L).

Sewage discharges	Observed BOD_5	Estimated <i>DOC</i>
Ciudad Guzmán	1,181	575
San Sebastián del Sur	1,363	663
Weighted concentration	1,272	620

Table 3 Diffuse sources of *SRP* in the Lake Zapotlan Basin.

Diffuse source	Hectares	<i>TP</i> (Kg/ha-yr)	<i>TP</i> (Kg/yr)	<i>SRP</i> (Kg/yr)
Irrigated agriculture	1,576	4.60	7,249.60	3,624.80
Temporary agriculture	17,425	2.00	3,485.00	17,425.00
Seeded pasture	2,834	0.10	283.40	141.70
Coniferous forest	11,960	0.11	1,315.60	657.80
Deciduous tropical forest	3,278	0.11	360.58	180.29
Secondary vegetation	3,020	0.10	302.00	151.00
Urban area	845	1.20	1,014.00	507.00
Lake and Wetlands	1,562	0.00	0.00	0.00
Atmospheric deposition over the entire basin	–	1.00	42,500.00	21,250.00
Total	42,500		87,875.18	43,937.59

Table 4 Diffuse sources of $NO_3 - N$ in the Lake Zapotlan Basin.


Diffuse source	Hectares	Exportation rate (Kg/ha-yr)	$NO_3 - N$ (Kg/yr)
Basin	42,500	2.43	103,275.00

Table 5 *TP* and *SRP* in Lake Zapotlan, in $\mu\text{gP/L}$.

Dry season (8 May 2003)		Rainy season (16 September 2003)	
Observed <i>TP</i>	Estimated <i>SRP</i>	Observed <i>TP</i>	Estimated <i>SRP</i>
916.00	549.60	1,122.00	673.20

The concentrations of $NO_3 - N$ analyzed according to official standard methods (CIATEJ [10]) were low (less than $50.00 \mu\text{gN/L}$) during both the dry and rainy seasons. Lewis [30] indicated that the denitrification process can restrict the supply of nitrogen to autotrophic organisms, and can eliminate all the nitrate in the water column in just a few days under anoxic conditions. These conditions could be occurring during the dry season in Lake Zapotlan, and therefore we assumed a $NO_3 - N$ concentration of $0 \mu\text{g/L}$ during this season. A $NO_3 - N$ concentration of $45 \mu\text{g/L}$ was assumed for the rainy season because of the load of nitrates to lake via surface. This value does not contradict the reported values by CIATEJ [10], and in this way the average annual $NO_3 - N$ concentration of the lake would be around $23.00 \mu\text{gN/L}$, a concentration of $NO_3 - N$ similar to that reported for the neighboring Lake Chapala [21], which has similar water quality of Lake Zapotlan.

The $NH_4 - N$ concentration was estimated from Figure 2 and Table 5. This figure is based on analysis of 450 lakes distributed around the world [41]. From Table 5, the measured *TP* concentration during the dry season in the lake was $916.00 \mu\text{gP/L}$. For this *TP* concentration, a $NO_3 - N : NH_4 - N$

Figure 2 *TP* versus $NO_3 - N : NH_4 - N$ rate [41].

ratio = 0.73 was provided by Figure 2. As the $NO_3 - N$ concentration of $0 \mu\text{gN/L}$ was assumed for the dry season, therefore the $NH_4 - N$ concentration is $0 \mu\text{gN/L}$. On the other hand, from Table 5, the measured *TP* concentration during the rainy season in the lake was $1,122.00 \mu\text{gP/L}$. For this *TP* concentration, a $NO_3 - N : NH_4 - N$ ratio = 0.70 was provided by Figure 2. As the $NO_3 - N$ concentration of $45 \mu\text{gN/L}$ was assumed for the rainy season, therefore the $NH_4 - N$ concentration is $64.30 \mu\text{gN/L}$.

3.5 TOC in the lake

Guzmán *et al.* [22] reported a mean *TOC* concentration in the lake of 11.50mgC/L . In the model, we used the assumption $DOC/POC = 6$ during the simulation period according to Wetzel [52]. Consequently, the *DOC* and *POC* mean concentration values can be calculated as 9.90mgC/L and 1.60mgC/L , respectively. *DOC* concentrations in freshwaters are generally in the range 2–10 mgC/L [48].

3.6 Un-ionized ammonia in the lake

Un-ionized ammonia nitrogen concentration n_g in the lake (in $\mu\text{gN/L}$) can be calculated as

$$n_g = \left(\frac{F_u}{1 - F_u} \right) n_a \quad (10)$$

where F_u = fraction of n_g in total ammonia nitrogen, which depends on temperature $T(^{\circ}\text{K})$ and pH [18],

$$F_u = \frac{1}{1 + 10^{0.09018 + \frac{2.729.92}{T} - pH}} \quad (11)$$

3.7 POM flux in the lake

We have considered that 50% of the BOD_5 in the lake is *POM* [7]. It is also possible to describe the flux of *POM* to the sediment (J_{C^*}) by using equivalents of oxygen ($\text{gO/m}^2 \cdot \text{d}$) [7],

$$J_{C^*} = v_p L_{p_w} \quad (12)$$

3.8 Nutrient flux in the lake

We have assumed that the nutrient concentration n ($\mu\text{g/L}$) in the lake depends linearly on the surface runoff, according to the equation:

$$n = n_E + \left(\frac{n_T - n_E}{Q_{2,T}} \right) Q_{2,k} \quad (13)$$

Flux F_n of nutrient n (in $\mu\text{g/m}^2 \cdot \text{d}$) at time t can be written as:

$$F_n = \frac{(n - \hat{n})(V \times 1000)}{A \times h} \quad (14)$$

We suspect that an important part of the flux of nutrients in the lake is eliminated from the water column and consumed by two main components of the ecosystem: aquatic weeds such as common cattail (*Typha latifolia*) and aquatic hyacinth (*Eichhornia crassipes*) and the harvest of fishes in the lake, mainly carp and tilapia. Unfortunately, one of the limitations of the proposed model is that it does not take into account a macrophyte and fishery compartment. Future work will further develop the model to take these components into consideration.

3.9 Total oxygen demand in the sediment

Total sediment oxygen demand (*TSOD*) (carbonaceous and nitrogenous, in $\text{gO/m}^2 \cdot \text{d}$) in the bottom sediments of the lake was calculated by applying the model developed by Di Toro *et al.* [16] that was solved by the secant modified method [8].

3.10 The ordinary differential equations system

The ordinary differential simultaneous equations (*ODE*) system describing the mass balance of different chemical and biological species describing the lake ecosystem (Table 6) were solved by using the fourth order Runge-Kutta method with a time increment of 0.1 days [8].

3.11 Model calibration

The solution of the *ODE*-system shown in Table 6 requires eight initial conditions. Through iterative simulations for the period of 2002–2003, we observed a cyclic annual pattern in the values of the state variables. Once we introduced a series of different initial conditions for the first year period, the starting conditions during the second year changed slightly and then remained almost constant. Hence, we established these conditions as the initial ones and the calculation process was performed in an iterative way until the *ODE*-system was stabilized. Finally, the stabilized initial conditions were considered as the initial conditions for the model. Table 7 shows the initial conditions and assumed nutrient loads to Lake Zapotlan starting on January 1st, 2002.

Nevertheless, it is possible that many different sets of initial conditions may allow the model to fit the observed data equally well yet yield different results in prediction. A numerical factorial experiment 2^5 was carried out within the model to estimate the susceptibility of the model to such behavior in terms of the *RMSE* error. With this purpose, the sets of initial conditions of the five state variables of Table 7 (ammonium nitrogen, nitrate nitrogen, *SRP*, particulate organic carbon, and dissolved organic carbon) were changed. We selected two levels for each initial condition of these mentioned variables. One level was fixed above the value showed in Table 7 and the other was fixed below this value, and consequently $2^5 = 32$ different sets of initial conditions were generated. After that, the model was run for each combination. During the calculation procedure, an *RMSE* error for each state variable was estimated from 32 combinations. The results and the fixed levels for each initial condition of the selected variables are shown in Table 8.

Once this step was accomplished, the goal of the calibration process was focused on the adjustment of the kinetics parameters until the model outputs reproduced the observed data in the monitoring stations, and also the reported water quality data observed

Table 6 Equations describing the model.

State variable	Symbol	Mass balance equation
Algae	a	$\frac{da}{dt} = \frac{Q_1 a_1}{V} + \frac{Q_2 a_2}{V} - \frac{Q_5 a}{V} + S_a$
Herbivorous zooplankton	z_h	$\frac{dz_h}{dt} = \frac{Q_1 z_{h,1}}{V} + \frac{Q_2 z_{h,2}}{V} - \frac{Q_5 z_h}{V} + S_{z_h}$
Carnivorous zooplankton	z_c	$\frac{dz_c}{dt} = \frac{Q_1 z_{c,1}}{V} + \frac{Q_2 z_{c,2}}{V} - \frac{Q_5 z_c}{V} + S_{z_c}$
Particulate organic carbon (<i>POC</i>)	c_p	$\frac{dc_p}{dt} = \frac{Q_1 c_{p,1}}{V} + \frac{Q_2 c_{p,2}}{V} - \frac{Q_5 c_p}{V} + S_{c_p}$
Dissolved organic carbon (<i>DOC</i>)	c_d	$\frac{dc_d}{dt} = \frac{Q_1 c_{d,1}}{V} + \frac{Q_2 c_{d,2}}{V} + \frac{Q_3 c_{d,3}}{V} - \frac{Q_5 c_d}{V} + S_{c_d}$
Ammonium nitrogen	n_a	$\frac{dn_a}{dt} = \frac{Q_1 n_{a,1}}{V} + \frac{Q_2 n_{a,2}}{V} - \frac{Q_5 n_a}{V} \pm \frac{F_{n_a} A}{V \times 1000} + S_{n_a}$
Nitrate nitrogen	n_i	$\frac{dn_i}{dt} = \frac{Q_1 n_{i,1}}{V} + \frac{Q_2 n_{i,2}}{V} - \frac{Q_5 n_i}{V} \pm \frac{F_{n_i} A}{V \times 1000} + S_{n_i}$
Soluble reactive phosphorus (<i>SRP</i>)	p_s	$\frac{dp_s}{dt} = \frac{Q_1 p_{s,1}}{V} + \frac{Q_2 p_{s,2}}{V} - \frac{Q_5 p_s}{V} \pm \frac{F_{p_s} A}{V \times 1000} + S_{p_s}$

Table 7 Initial conditions and nutrient load to Lake Zapotlan.

State variable	Symbol	Units	Sewage discharges	Surface runoff	Initial condition
Herbivorous zooplankton	z_h	mgC/L	0	0	0
Ammonium nitrogen	n_a	$\mu\text{gN/L}$	0	0	1.27
Nitrate nitrogen	n_i	$\mu\text{gN/L}$	Eq. (5)	5,461.77 (Eq. (9))	0.90
Soluble reactive phosphorus	p_s	$\mu\text{gP/L}$	Eq. (6)	2,323.67 (Eq. (8))	551.53
Algae	a	$\mu\text{gChla/L}$	1	5	4.39
Carnivorous zooplankton	z_c	mgC/L	0	0	0.25
Particulate organic carbon	c_p	mgC/L	350	8.8 ($c_p = 0.1c_d$ [52])	0.61
Dissolved organic carbon	c_d	mgC/L	620 (Table 2)	88	5.23

Table 8 Levels and RMSE of the state variables.

State variables	z_h	n_a	n_i	P_s	a	Z_c	c_p	c_d
Level (lower, upper)	–	(1, 2)	(0, 1)	(500, 600)	–	–	(0.5, 1)	(5, 10)
RMSE error	0.00	0.03	0.00	0.01	0.04	0.00	0.00	0.06

by Guzmán *et al.* [22] for Lake Zapotlan, and by Wetzel [52] for other lakes.

4 Results and discussion

The concentrations of the inorganic nutrients in the lake were positively correlated with the lake temperature, lake inflow, pH , DO , $TSOD$, herbivorous and carnivorous zooplankton concentration, nutrients flux and TOC in the lake, and negatively correlated with the lake outflow and POM flux in the lake ($P < 0.0001$). There was no correlation between herbivorous zooplankton concentrations in the lake with other variables, but there was a high positive correlation between carnivorous zooplankton and lake temperature, photoperiod, lake inflow, pH , DO , $TSOD$, ammonium flux, nutrient concentration, algal concentrations and TOC ($P < 0.0001$). Algal concentrations were positively correlated with the lake temperature, photoperiod, lake inflow, pH , DO , $TSOD$, ammonium flux, nutrient concentration, carnivorous zooplankton and TOC concentration; and negatively correlated with lake outflow and POM flux ($P < 0.0001$). $TSOD$ was positively correlated with the lake temperature, lake inflow, pH and DO , nutrient concentration, ammonium flux, and herbivorous and carnivorous zooplankton concentrations and TOC in the lake; and was negatively correlated with the lake outflow and POM flux in the lake ($P < 0.0001$). Lake residence time Θ was positively correlated with the depth; and negatively correlated with the lake outflow and solar radiation ($P < 0.0001$). From Figure 3 we assumed that nitrogen was the limiting nutrient in the water body, since SRP simulated values were always in excess of total N values during the simulation period. This figure shows that simulated N and P values peaked in July to September, with minima in the winter (December to April).




Figure 3 Simulation of the inorganic nutrients in Lake Zapotlan.

In terms of nutrient fluxes (Figure 4) we observe a minimum in the fluxes of nitrates, ammonium, and SRP in March to May (spring) with a subsequent peak in fluxes in the summer, corresponding to the peak in nutrient concentrations. The March–April–May minimum in the downward flux of nutrients is due to the intensity of the upward flux of nutrients (mainly caused by resuspension) because of decreased water level in spring and correlated with the monthly average wind speed. During this quarter the strongest winds blow across Lake Zapotlan (see Figure 5). The increase in nutrient concentrations in the lake corresponds to an increase in algal biomass, with a subsequent increase in biomass of herbivorous and carnivorous zooplankton (see Figure 6).

During the rainy season, we observed an increase in the nutrient concentrations in the lake, and consequently pH , DO ,




Figure 4 Simulation of the nutrient fluxes in Lake Zapotlan.




Figure 5 Basic atmospheric parameters for Lake Zapotlan.

TSOD, flux of nutrients, and *DOC* increased as well. On the other hand, the flux of *POM* to the bottom sediments decreased. Using Eq. (11), we calculated that in the rainy season more than 62% of the total ammonia was in the form of gaseous ammonia, whose maximum concentration exceeded 100 $\mu\text{gN/L}$, a concentration level that is normally toxic for most fish species. Using the observed monitoring data, the $\text{NO}_3 - \text{N}$ concentration in the model was always less than 50 $\mu\text{g/L}$ during the simulated period and the observed *SRP* concentration was adjusted to the model (see Figure 3). During the rainy season there was an increased flux of nutrients from the water column to the bottom sediments (Figure 4). Therefore the simulation shows an increase in the values of *pH*, *DO*, *TSOD*, nutrient concentrations in the water column and *DOC*. During the dry season, the flux of ammonium and *SRP* from the water column to the bottom sediments decreased, and nitrates entered the water column from bottom sediments.

Figure 6 shows the results of the food chain simulation in the lake. Zooplankton concentrations were expressed in chlorophyll-*a* concentration units in order to compare them to the behavior of algal concentrations [7]. Results show strong interaction between predators and prey. Algae showed a rapid increase in

concentration during the second quarter of the year, and then decreased gradually until December. The maximum herbivorous zooplankton concentration level occurred in May and the minimum occurred in February.

Figure 7 shows the difference in the productivity between the rainy season and the dry season. High organic carbon generation during the rainy season has several implications. Its decomposition can have an impact on the oxygen content of bottom waters and the aquatic biota. Besides, the transport and fate of substances such as heavy metals can be strongly associated with organic matter [7]. At the same time that organic carbon generation increases, the flux of *POM* to the bottom sediments decreases, while *pH*, *DO*, *TSOD*, the flux of ammonium from the water column, nutrients, herbivorous and carnivorous zooplankton concentrations in the lake increase. As a result of the simulation, we obtained a *DOC/POC* rate of 6 [52] and the same *TOC* mean concentration reported by Guzmán *et al.* [22]. Figure 8 shows the *TSOD* and the flux of *POM* to the bottom sediments of the lake. Maximum of *TSOD* occurred during the rainy season, when minimum fluxes of *POM* to the bottom sediments occurred.




Figure 6 Food chain simulation in Lake Zapotlan.




Figure 7 Organic carbon simulation in Lake Zapotlan.




Figure 8 Sediment oxygen demand and flux of POM in Lake Zapotlan.

5 Conclusions

Lake Zapotlan is a dynamic ecosystem, dominated by the interactions amongst solar radiation, surface runoff and discharge of wastewater. In early spring, when the wind restricts the flux of SRP and ammonia to the bottom and mixes nitrate throughout the water column, we have observed an increase in primary productivity in the lake. The calculated increase in primary productivity is attributable to the diffuse loading of nutrients mainly during the rainy season, and to the slow increase of the air temperature, nitrification, hydrolysis of DOC, carnivorous zooplankton respiration, etc. Primary productivity decreases at the beginning of the summer because of the following factors:

- Increase in the ammonium flux to the bottom sediments;
- diminution of the nitrate flux from the bottom sediments to the water column;
- increase of the POM flux to the bottom sediments; and the
- limited amount of TOC present in the water column.

The lowest levels of nutrients and primary production occur in winter and at the beginning of spring, principally due to low levels of DOC in the lake, which results in a low generation of nutrients via hydrolysis.

The simplified nutrients/food chain dynamic model developed in this work permits a better understanding of the processes occurring in the lake. It describes the impact of the point and non-point sources of nutrients on the water quality of the lake and the growth and composition of the food chain. The main point source of nutrients affecting the water quality in the system is the direct discharge of poorly treated municipal sewage into the lake. It is necessary to improve sewage treatment by removing nutrients prior to its discharge into the lake. Alternatively one could consider diversion of treated sewage to cultivated fields (reuse of water). This measure would also minimize the extraction of groundwater from wells. The control of the diffuse source of nutrients is a complicated issue that requires the development of a management plan for the entire lake basin. A rapid action plan that minimizes the load of nutrients from agricultural areas

surrounding the lake could involve the creation of a vegetated, non-cultivated riparian zone around the lake, acting as a buffer to capture the excess of nutrients. Presently there is no system protecting the shoreline of the lake from the agrochemicals used in local agriculture.

The proposed model integrates the hydrologic balance, the energy equations, and the food chain processes in the lake. Therefore, it is possible to simulate different scenarios to improve the water quality of the system by reducing the loads of nutrients. This tool permits decision makers to focus administrative and economic resources on those measures that give better results in terms of water quantity and quality in the lake.

The next step in the development of this model will be to include a fisheries and aquatic macrophytes component, as noted earlier. Another of the areas of concern in this lake is the existence of blue-green algae blooms producing cyanotoxins. This model will be improved to determine the processes that permit the development of such algae.

Notation

- a = Concentration of algae in the Lake ($\mu\text{gChla/L}$)
- A = Area of the sediment-water interface \approx Lake area (m^2)
- a_j = Concentration of algae in the flow j ($\mu\text{gChla/L}$)
- BOD_5 = 5-day biochemical oxygen demand
- c = Concentration of state variable v in the lake ($\mu\text{g/m}^3$ or mg/m^3)
- c_j = Concentration of state variable v in the flow i ($\mu\text{g/m}^3$ or mg/m^3)
- c_d = Concentration of DOC in the Lake (mgC/L)
- $c_{d,j}$ = Concentration of DOC in the flow j (mgC/L)
- c_p = Concentration of POC in the Lake (mgC/L)
- $c_{p,j}$ = Concentration of POC in the flow j (mgC/L)
- DOC = Dissolved organic carbon
- f = Photoperiod
- F_n = Flux of nutrient n at time t ($\mu\text{g/m}^2 \cdot \text{d}$)
- F_v = Flux of the state variable v in the interface (when nutrient) ($\mu\text{g/m}^2 \cdot \text{d}$) (A positive sign denotes a flux directed towards the water column, a negative sign denotes a opposite flux).
- h = Time increment (days)
- I_0 = Solar radiation at the lake surface
- j = Flow = 1(sewage), 2(runoff), 3(surface infiltration), 4(precipitation), 5(water extraction)
- J_{C^*} = flux of POM to the sediment ($\text{gO/m}^2 \cdot \text{d}$)
- $k(T)$ = process velocity at temperature T
- k_{20} = process velocity at 20°C
- L_{pw} = POM concentration in the lake in the time t (mgO/L)
- n_a = Concentration of $\text{NH}_4 - \text{N}$ in the Lake ($\mu\text{gN/L}$)
- $n_{a,j}$ = Concentration of $\text{NH}_4 - \text{N}$ in the flow j ($\mu\text{gN/L}$)
- n_E = Concentration during the dry season ($\mu\text{g/L}$)
- n_i = Concentration of $\text{NH}_4 - \text{N}$ in the Lake ($\mu\text{gN/L}$)

n_g = Concentration of un-ionized $NH_3 - N$ in the Lake ($\mu\text{gN/L}$)
 $n'_{i,1}$ = Concentration of $NO_3 - N$ at the sewage on May 8, 2003 ($\mu\text{gN/L}$)
 $n_{i,j}$ = Concentration of $NH_4 - N$ in the flow j ($\mu\text{gN/L}$)
 n_T = Concentration in the month of maximum surface runoff ($\mu\text{g/L}$)
 \hat{n} = Concentration of nutrient n calculated by the model without considering any flux ($\mu\text{g/L}$)
 DO = Concentration of oxygen dissolved in the Lake ($\mu\text{g/L}$)
 POC = Particulate organic carbon
 POM = Particulate organic matter
 p_s = Concentration of SRP in the Lake ($\mu\text{gP/L}$)
 $p_{s,j}$ = Concentration of SRP in the flow j ($\mu\text{gP/L}$)
 $p'_{s,1}$ = Concentration of SRP at the sewage on May 8, 2003 ($\mu\text{gP/L}$)
 Q_j = Volumetric flow rate of the flow j (m^3/d)
 $Q_{2,k}$ = Surface runoff in the month k (m^3/month)
 $Q_{2,T}$ = Maximum surface runoff during the rainy season (m^3/month)
 \bar{Q}_2 = Annual average of surface runoff (L)
 Q'_1 = Sewage discharges to the lake on May 8, 2003 (m^3/d)
 S_v = Sources and sinks of the state variable v ($\mu\text{g/L} \cdot \text{d}$), calculated by the Chapra model [7].
 $RMSE$ = Root mean square error
 SRP = Soluble reactive phosphorus
 t = Time (d)
 T = Lake temperature ($^{\circ}\text{C}$)
 TOC = Total organic carbon
 TP = Total phosphorus
 $TSOD$ = Total oxygen demand in the bottom sediments of the lake ($\text{gO}/\text{m}^2 \cdot \text{d}$)
 v = State variable = $a, z_h, z_c, c_p, c_d, n_a, p_s$ or n_i
 V = Lake volume (m^3)
 v_p = POM settlement velocity = 0.2 m/d [7]
 $W'_{n_i,1}$ = Mass loads of $NO_3 - N$ from point sources on May 8, 2003 (μg)
 $W'_{p_s,1}$ = Mass load of SRP from point sources on May 8, 2003 (μg)
 \bar{W}'_{n_i} = Annual diffuse loads of $NO_3 - N$ from the basin (μg)
 \bar{W}'_{p_s} = Annual diffuse loads of SRP from the basin (μg)
 z_c = Concentration of carnivorous zooplankton in the Lake (mgC/L)
 $z_{c,j}$ = Concentration of carnivorous zooplankton in the flow j (mgC/L)
 z_h = Concentration of herbivorous zooplankton in the Lake (mgC/L)
 $z_{h,j}$ = Concentration of herbivorous zooplankton in the flow j (mgC/L)
 θ = Temperature factor
 Θ = Lake residence time (months)

References

1. AMBROSE, R.B., WOOL, T.A. and MARTIN, J.L. (1993). The Water Quality Analysis Simulation Program, WASP5. Part A: Model documentation. USEPA ERL, Athens, GA 30613.
2. ARHONDITSIS, G.B. and BRETT, M.T. (2005). "Eutrophication Model for Lake Washington (USA)," *Ecological Modelling*, 187(2–3), 179–200.
3. BOWIE, G.L., MILLS, W.B., PORCELLA, D.B., CAMPBELL, C.L., PAGENKOPF, J.R., RUPP, G.L., JOHNSON, K.M., CHAN, P.W., GHERINI, S.A. and CHAMBERLAIN, C.E. (1985). Rates, Constants, and Kinetic Formulation in Surface Quality Modeling. U.S.E.P.A., ORD, Athens, GA, ERL, EPA/600/3-85/040.
4. CANALE, R.P., HINEMANN, D.F. and NACHIPPAN, S. (1974). A Biological Production Model for Grand Traverse Bay. University of Michigan Sea Grant Program, Technical Report No. 37, Ann Arbor MI.
5. CEASJ (2003). Plan Maestro de la Laguna de Zapotlán, Consejo Estatal de Agua y Saneamiento del Gobierno del Estado de Jalisco, Ciudad Guzmán, México. 111 pp.
6. CHAMBERS, P.A. and DALE, A.R. (1996). Contribution of Industrial, Municipal, Agricultural and Groundwater Sources to Water Quality in the Athabasca and Wapiti-Smoky Rivers. Northern River Basins Study Technical Report No. 110, Alberta, Canada.
7. CHAPRA, S.C. (1997). Surface Water-Quality Modeling, WCB/McGraw-Hill, New York, 844 pp.
8. CHAPRA, S.C. and CANALE, R.P. (1988). Numerical Methods for Engineers, 2nd Edn., McGraw-Hill, New York, 812 pp.
9. CHEN, C.W. and ORLOB, G.T. (1975). Ecological simulation for aquatic environments, in B.C. PATTON (ed.), Systems Analysis and Simulation in Ecology, Vol. III, Academic, New York, 475 pp.
10. CIATEJ. (2003). Resultados analíticos de calidad de agua del Lago de Zapotlán, Laboratorio de Química Analítica y Metrología, Guadalajara, Jalisco.
11. CNA (2004). Meteorological monthly reports. National Water Commission. Department of Hydrometric. Meteorological station of Ciudad Guzmán. Mexico.
12. COLE, T.M. and BUCHAK, E.M. (1995). CE-QUAL-W2: A Two-Dimensional, Laterally Average, Hydrodynamic and Water Quality Model, Version 2. U.S. Army Corps of Engineers, Waterways Experiment Station, Tech. Report EL-95-x.
13. DE ANDA, J. (2000). Contaminación por Fósforo en el Lago de Chapala, Ph.D. Thesis, Unidad de Ciclos Profesionales y de Posgrado, Colegio de Ciencias y Humanidades, UNAM, 122 pp. México, D.F.
14. DE ANDA, J., SHEAR, H., MANIAK, U. and RIEDEL, G. (2001). "Phosphates in Lake Chapala, México," *Lakes & Reservoirs: Research and Management*, 6, 313–321.
15. DI TORO, D.M. and MATYSTIK, W.F. (1979). "Phosphorus Recycle and Chlorophyll in the Great Lakes," *J. Great Lakes Research*, 5, 233–245.

16. DI TORO, D.M., PAQUIN, P.R., SUBBURAMU, K. and GRUBER, D.A. (1990). "Sediment Oxygen Demand Model: Methane and Ammonia Oxidation," *J. Environ. Engr. Div. ASCE*, 116(5), 945–986.
17. DODDS, W.K. (2003). "Misuse of Inorganic N and Soluble Reactive P Concentrations to Indicate Nutrient Status of Surface Waters," *J. N. Am. Benthol. Soc.*, 22(2), 171–181.
18. EMERSON, K., RUSSO, R.C., LUND, R.E. and THURSTON, R.V. (1975). "Aqueous Ammonia Equilibrium Calculations: Effect of pH and Temperature," *Journal of the Fisheries Research Board of Canada*, 32, 2379–2383.
19. EPPLEY, R.W. (1972). "Temperature and Phytoplankton Growth in the Sea," *Fishery Bulletin*, 70(4), 1063–1085.
20. FADINI, P.S., JARDIM, W.F. and GUIMARÃES, J.R. (2004). "Evaluation of Organic Load Measurement Techniques in a Sewage and Waste Stabilization Pond," *J. Braz. Chem. Soc.*, Vol.15 No.1 São Paulo Jan./Feb. (2004).
21. FAO (Food and Agricultura Organization) (2000). Reporte de la Iniciativa de la Ganadería, el Medio Ambiente y el Desarrollo (LEAD) – Integración por Zonas de la Ganadería y de la Agricultura Especializada (AWI) – Opciones para el Manejo de Efluentes de Granjas Porcícolas de la Zona Centro de México, INIFAP, UNAM y Swiss College of Agricultural.
22. GUZMÁN, M., MICHEL, J.G., PEREDO, A., GARCÍA, M. and CAMARENA, S. (2000). Limnología de la Laguna de Zapotlán, CUCBA, Reporte interno Universidad de Guadalajara. Ciudad Guzmán, México.
23. HAMBLIN, P.F., BOOTSMA, H.A. and HECKY, R.E. (2003). "Modeling Nutrient Upwelling in Lake Malawi/Nyasa," *Journal of Great Lakes Research*, 29, 34–47.
24. HydroQual. (2001). Analysis of the Addition of a Third Algal Group to the Bays Eutrophication Model (BEM) kinetics. Boston: Massachusetts Water Resources Authority. Report 2001–15. 110 pp.
25. INEGI. (2000). XII Censo General de Población y Vivienda, México, D.F.
26. JANSE, J.H. (1997). "A Model of Nutrient Dynamics in Shallow Lakes in Relation to Multiple Stable States," *Hydrobiologia*, 342–343(8), 1–8.
27. JANSE, J.H., VAN DONK, E. and ALDENBERG, T. (1998). "A Model Study on the Stability of the Macrophyte-Dominated State as Affected by Biological Factors," *Water Research*, 32(9), 2696–2706.
28. JANSE, J.H., VAN DONK, E. and GULATI, R.D. (1995). "Modeling Nutrient Cycles in Relation to Food Web Structure in a Biomanipulated Shallow Lake," *Neth. J. Aquat. Ecol.*, 29(1), 67–79.
29. LAWS, E.A. and CHALUP, M.S. (1990). "A Microalgal Growth Model," *Limnol. Oceanogr.*, 35(3), 597–608.
30. LEWIS, W.M. (2002). "Causes for the High Frequency of Nitrogen Limitation in Tropical Lakes," *Verh. Internat. Verein. Limnol.*, 28, 210–213.
31. LEWIS, W.M., MELACK, J.H., MCDOWELL, W.H., MCCLAIN, M. and RICHEY, J.E. (1999). "Nitrogen Yields from Undisturbed Watersheds in the Americas," *Biogeochemistry*, 46, 149–162.
32. MCFARLAND, A. and HAUCK, L. (1998). Lake Waco/Bosque River Watershed Initiative Report: Determining Nutrient Contribution by Land Use for the Upper North Bosque River Watershed. TIAER PR 98-01.
33. Minuteman Software. 2000. *GPSS World Tutorial Manual*. Holly Springs, NC, U.S.A.
34. MOURAD, D.S.J., VAN DER PERK, M., GOOCH, G.D., LOIGU, E., PIIRIMAE, K. and STALNACKE, P. (2005). "GIS-Based Quantification of Future Nutrient Loads into Lake Peipsi/Chudskoe Using Qualitative Regional Development Scenarios," *Water Science and Technology*, 51(3–4), 355–363.
35. NURNBERG, G.K. and LAZERTE, B.D. (2004). "Modeling the Effect of Development on Internal Phosphorus Load in Nutrient-Poor Lakes," *Water Resources Research*, 40(1), 1051–1059.
36. ORTIZ-JIMÉNEZ, M.A., DE ANDA, J. and SHEAR, H. (2005). "Hydrologic Balance of Lake Zapotlán, México," *Journal of Environmental Hydrology*, 13(5).
37. OSIDELE, O.O. and BECK, M.B. (2004). "Food Web Modelling for Investigating Ecosystem Behaviour in Large Reservoirs of the South-Eastern United States: Lessons from Lake Lanier, Georgia," *Ecological Modelling*, 173(2–3), 129–158.
38. PORTIELJE, R. and RIJSDIJK, R.E. (2003). "Stochastic Modelling of Nutrient Loading and Lake Ecosystem Response in Relation to Submerged Macrophytes and Benthivorous Fish," *Freshwater biology*, 48(4), 741–755.
39. PRASUHN, V. and SIEBER, U. (2005). "Changes in diffuse phosphorus and nitrogen inputs into surface waters in the Rhine watershed in Switzerland," *Aquatic Sciences*, 67(3), 363–371.
40. PUIJENBROEK, P.J., JANSE, J.H. and KNOOP, J.M. (2004). "Integrated Modelling for Nutrient Loading and Ecology of Lakes in The Netherlands," *Ecological Modelling*, 174(1–2), 127–141.
41. QUIRÓS, R. (2003). "The relationship between Nitrate and Ammonium Concentration in the Pelagic Zone of Lakes," *Limnetica*, 22(1–2), 37–50.
42. RANDALL, G.W. and MULLA, D.J. (2001). "Nitrate Nitrogen in Surface Waters as Influenced by Climatic Conditions and Agricultural Practices," *J. Environ. Qual.*, 30, 337–344.
43. RILEY, G.A. (1956). "Oceanography of Long Island Sound 1952–1954, II. Physical Oceanography," *Bull. Bingham Oceanog. Collection*, 15, 15–16.
44. RULEY, J.E. and RUSCH, K.A. (2005). "Development of a Simplified Phosphorus Management Model for a Shallow, Subtropical, Urban Hypereutrophic Lake," *Ecological Engineering*, 22(2), 77–98.
45. RYDING, S.O. and RAST, W. (eds.) (1989). The Control of Eutrophication of Lakes and Reservoirs. UNESCO, Man And The Biosphere Series. Vol. 1. The Parthenon Publishing Group. 314 pp.

46. SCHNOOR, J. (1996). *Environmental Modeling*. Wiley-Interscience. New York, 682 pp.
47. SUDARSHAN, N.T. (1995). *Planktonic Metabolism in Lake Marion, SC: Functional Response to Light, Temperature, and Nutrient Concentrations*. Masters Thesis, Department of Environmental Health Sciences, University of South Carolina.
48. TALLING, J.F. and LEMOALLE, J. (1998). *Ecological Dynamics of Tropical Inland Waters*. Cambridge University Press, Cambridge, MA. 441 pp.
49. THOMANN, R.V. and MUELLER, J.A. (1987). *Principles of Surface Water Quality Modeling*, Harper Collins Publishers, Inc., 644 pp.
50. USEPA. (2000). *Nutrient Criteria Technical Guidance Manual. Lakes and Reservoirs*. First Edition. United States Office of Water EPA-822-B00-001, Washington, DC 20460.
51. WANG, S.H., HUGGINS, D.G., FREES, L., VOLKMAN, C.G., LIM, N.C., BAKER, D.S., SMITH, V., and DEHOYELLES, F. (2005). "An Integrated Modeling Approach to Total Watershed Management: Water Quality and Watershed Assessment of Cheney Reservoir, Kansas, USA," *Water Air and Soil Pollution*, 164(1-4), 1-19.
52. WETZEL, R.G. (2001). *Limnology. Lake and River Ecosystems*. 3rd edn., Academic Press, San Diego, 1006 pp.
53. ZHANG, J. and JORGENSEN, S.E. (2005). "Modelling of Point and Non-Point Nutrient Loadings from a Watershed," *Environmental Modelling and Software*, 20(5), 561-574.

