

## A Note on Serum Insulin in Mexican Cuino Pigs

<sup>1</sup>F. Grageola, <sup>1</sup>C. Lemus, <sup>2</sup>R.A. Alonso, <sup>2</sup>C. Camacho and <sup>3</sup>J. Ly

<sup>1</sup>Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Nayarit  
Ciudad Universitaria, Amado Nervo, Tepic. Nayarit, México

<sup>2</sup>Facultad de Medicina Veterinaria y Zootecnia, Universidad Nacional Autónoma de México, México,  
D.F. Ciudad Universitaria, Circuito Exterior, México, D.F. 04510, México

<sup>3</sup>Instituto de Investigaciones Porcinas, P.O. Box 1, Punta Brava, La Habana, Cuba

**Abstract:** In this study a 2×2 factorial arrangement was used to study the effect of sex and age on serum insulin levels in 24 growing Mexican Cuino pigs between 3 and 6 months old. There were no significant differences ( $p < 0.05$ ) neither for sex × age interaction nor for sex in any measured index. Overall, Mexican Cuino pigs had 14.4 and 50.8 kg of live weigh ( $p < 0.001$ ) and 9.33 and 19.26 u $\mu$ L<sup>-1</sup> of fasting serum insulin levels ( $p < 0.001$ ) at 3 and 6 months old, respectively. Pearson correlation matrix revealed significant differences ( $p < 0.05$ ) among age, live weigh and fasting insulin levels in the examined period of life in growing Mexican Cuino pigs. According to the present evaluation, hiperinsulinemia should be present in Mexican Cuino pigs, as it has been found in either obese conventional genotypes of pigs or genetically manipulated to be small and obese animals. This effect was accentuated as pigs aged up to 6 months life. Sex appeared to have no influence on serum status of Mexican Cuino pig insulin.

**Key words:** Mexican cuino pigs, insulin, age, body weight, sex

### INTRODUCTION

Cuino pigs belong to one of three local Mexican genotypes which have been adapted to the habitat after near 5 centuries of introduction from East Asia (Lemus and Alonso, 2005). This type of animal is characterized for attaining the adult stage when arriving to approximately 50 kg. In this moment, the Mexican Cuino pigs have a height of 53-65 cm on average, since they have a small body with a marked trend to accumulate fat (Lemus *et al.*, 2003, 2005). In this connection, it has been suggested that adiposity in animals such as pigs is related to some extent, to the mechanism of insulin control of metabolism (Elsaesser *et al.*, 2002). In this sense, it has been argued that insulin is a crucial regulator of lipids, through different actions and among them, the stimulation of preadipocyte differentiation to adipocytes, glucose cell transport mechanism, triglyceride synthesis (Walton and Etherton, 1986) and lipolysis inhibition (Dunshea *et al.*, 1992b). On the other hand, it has been claimed that antilipolytic hormones as insulin, have a major effect on subcutaneous than visceral lipid tissue.

In our knowledge, there is not previous information related to the role played by insulin in Mexican Cuino pigs, although insulin status has been examined in other

Mexican genotypes employed for studying carbohydrate metabolism (Phillips *et al.*, 1979, 1982). In this respect, the Mexican Cuino genotype of pigs could be used in metabolic studies concerning obesity, as it has been done with other local, Mexican breed of small size, due to be well suited for using at laboratory scale, among other reasons (Panepinto *et al.*, 1978).

The aim of the present study was the estimation of serum levels of insulin in Mexican Cuino pigs for increasing the knowledge of its physiology status, as attained in natural conditions of animal husbandry.

**Area of the study:** The study was conducted in the Faculty of Veterinary Medicine and Zootechnics of the Autonomous University of Nayarit, at Tepic, Mexico. The climate was semi-warmth and sub-humid, with rainy summers and average annual temperature of 22°C.

### MATERIALS AND METHODS

**Animals and diets:** A total of growing 24 Mexican pigs of the local, Cuino genotype, from the Faculty herd were used. There were 12 animals of three and another of six months old, castrate male and female in the same proportion. The animals were given ad libitum a

commercial balanced ration containing either 16 or 14% crude protein (N×6.25) for animals growing up to 3 and 6 months of age, respectively. Housing consisted of pens in an open stable. Management of animals was conducted as previously described (Grageola, 2007).

**Sampling and analysis:** Live weight of pigs was determined in the day of blood extraction. Blood sampling was carried out in the morning after a 24 h fasting period. Two different methods for blood extraction were employed. In young, 3 months old pigs blood was obtained from vena cava, whereas the orbital sinus (Morton *et al.*, 1993) was used for blood sampling in sixth months old pigs. Blood was collected by using plastic syringes and immediately transferred to iced test tubes for transporting to the laboratory. Afterwards the serum was obtained by blood centrifugation at 5 000 rpm during 5 min. Pig serum was stored at -20°C until analysis. Serum samples were analyzed by duplicate according to a radioimmunoassay by using a <sup>125</sup>I-RIA commercial kit. Pig serum samples were prepared following the instructions of the manufactures, according to conventional procedures (Hales and Randle, 1963; Steele *et al.*, 1985), with porcine insulin as standard.

**Statistical analysis:** Statistical analysis were performed using ANOVA techniques of SAS (1987). Means were compared by least significant differences at p<0.05, according to a GLM procedure. Factors in the model included sex, age and sex × age interaction, whereas serum insulin and body weight were response variables.

**RESULTS AND DISCUSSION**

There were no significant differences (p>0.05) for neither sex × age interaction nor for sex in any measured index. Therefore, data were pooled regardless of the sex effect. Overall, Mexican Cuino pigs had 14.4 and 50.8 kg live weight (p<0.001) at 3 and 6 months old, respectively (Table 1) and this was in agreement of the slow growth rate of this genotype, as to be expected (Lemus and Alonso, 2005).

A significant effect (p<0.001) was encountered for fasting serum insulin levels when data of animals of 3 and 6 months old were contrasted (9.33 and 19.26 u mL<sup>-1</sup>, respectively). In this connection, blood insulin levels has been found in pigs to be affected by several factors, not only from the nutritional status point of view (Atinmo *et al.*, 1976; Buonomo and Baile, 1991), but also by other causes such as physiological conditions (Steele *et al.*, 1985; Dunshea *et al.*, 1992a, b, c) and breed (Wangsness *et al.*, 1977). Even more, Elsaesser *et al.*

Table 1: Effect of age on live weight and serum insulin in Mexican Cuino pigs

	Age, months		SE±
	3	6	
n	12	12	-
Live weight, kg	14.4	50.8	1.4***
Serum insulin, u mL <sup>-1</sup>	9.33	19.26	2.82***

\*\*\* p<0.001

Table 2: Pearson correlation matrix of growth and insulin indices in Mexican Cuino pigs (n = 24)

	Age	Live weight
Live weight	0.982***	0.585***
Insulin	0.559**	

\*\* p<0.01; \*\*\* p<0.001

(2002) obtained evidences that divergent pig genotypes, the small, obese Gottingen miniature and the large, lean German Landrace animal could reflect differences in blood insulin status. In fact, according to Elsaesser *et al.* (2002), obese pigs exhibited fasting plasma insulin values evidently higher than that of the Landrace breed. In the particular case of lean and genetically obese pigs, Wangsness *et al.* (1977) observed that at 1, 3 and 6 months of age, obese pigs were not clearly hyperinsulinemic animals but had lower plasma somatotropin than lean pigs.

As it is shown in Table 2 data from the Pearson Matrix Correlation revealed a clear, strong interdependence (p<0.001) was observed between age and sex of young Mexican Cuino pigs. In this connection, serum insulin values were also significantly correlated (p<0.01) to either age or sex of the animals, although the coefficient of determination did not reached 0.6 values. According to these findings, it could be assumed that fasting insulin response to age in young Mexican Cuino pigs could be somewhat linear up to six months of age, according to the following expression

$$y = 5.39 + 0.07 x, SD \pm 0.02$$

Where, y and x expressed fasting insulin levels (in U mL<sup>-1</sup>) and age (in days), respectively.

In summary, according to the present evaluation, hiperinsulinemia should be present in Mexican Cuino pigs, as it has been found, with the exception of Weiler *et al.* (1998), in obese conventional genotypes of pigs (McCuskers *et al.*, 1985) or in genetically manipulated to be small and obese animals (Wangsness *et al.*, 1981; Mersmann *et al.*, 1982; Elsaesser *et al.*, 2002). This effect was accentuated as pigs aged up to six months life. Sex appeared to have no influence on serum status of Mexican Cuino pig insulin.

**REFERENCES**

- Atinmo, T., C. Baldijao, W.G. Pond and R.H. Barnes, 1976. Plasma insulin levels in weaned pigs fed protein or energy restricted diets. *J. Nutr.*, 106: 1654-1658.
- Buonomo, F.C. and C.A. Baile, 1991. Influence of nutritional deprivation on insulin-like growth factor I, somatotropin and metabolic hormones in swine. *J. Anim. Sci.*, 69: 755-760.
- Dunshea, F.R., D.E. Bauman, R.D. Boyd and A.W. Bell, 1992a. Temporal response of circulating metabolites and hormones during somatotropin treatment of growing pigs. *J. Anim. Sci.*, 70: 123-131.
- Dunshea, F.R., D.M. Harris, D.E. Bauman, R.D. Boyd and A.W. Bell, 1992b. Effect of somatotropin and glycerol metabolism in growing pigs. *J. Anim. Sci.*, 70: 132-140.
- Dunshea, F.R., D.M. Harris, D.E. Bauman, R.D. Boyd and A.W. Bell, 1992c. Effect of porcine somatotropin on *in vivo* glucose kinetics and lipogenesis in growing pigs. *J. Anim. Sci.*, 70: 141-151.
- Elsaesser, F., M.W. Pfaffl, H.D.H. Meyer, B. Serpek and H. Sauervein, 2002. Differences in the somatotropic axis in blood cortisol, insulin and thyroid hormone concentrations between 2 pig genotypes with markedly divergent growth rates and the effects of growth hormone treatment. *Anim. Sci.*, 74: 423-430.
- Grageola, F., 2007. Caracterización de la hormona sérica insulina en el estudio de la obesidad del cerdo Cuino. Tesis MVZ. Univ. Autón. Nayarit. Tepic, pp: 24.
- Hales, C.N. and P.J. Randle, 1963. Immunoassay of insulin with insulin antibody precipitate. *Biochem. J.*, 88: 137-146.
- Lemus, C. and M.L. Alonso, 2005. El Cerdo Pelón Mexicano y otros Cerdos Criollos. Editorial de la Universidad Autónoma de Nayarit. Tepic, pp: 251.
- Lemus, C., R. Alonso, J.G. Herrera, M.L. Alonso, N.R. Ramírez and D. Mota, 2005. Growth, morphometry and reproductive performance of creole Cuino pigs in Mexico. *J. Anim. Vet. Adv.*, 4: 855-858.
- Lemus, C., R. Ulloa, M. Ramos, F.J. Estrada and N.R. Ramírez, 2003. Morphologic characteristics in Mexican native pig. *Arch. Zootecnia*, 52: 105-108.
- McCusker, R.H., P.J. Wangsness, L.C. Griel and J.F. Kavanaugh, 1985. Effects of feeding, fasting and refeeding on growth hormone and insulin in obese pigs. *Physiol. Behav.*, 35: 383-388.
- Mersmann, H.J., W.G. Pond and J.T. Yen, 1982. Plasma glucose, insulin and lipids during growth of genetically lean and obese swine. *Growth*, 46: 189-198.
- Morton, D.B., D. Abbot, R. Barclay, B.S. Close, R. Ewband and D. Gask *et al.*, 1993. Removal of blood from laboratory mammals and birds: First Report of the BVA/FRAME/RSPCA/UFAW Joint Working Group of Refinement. *Lab. Anim.*, 27: 1-22.
- Panepinto, L.M., R.W. Phillips, R.W. Wheeler and D.H. Will, 1978. The Yucatan miniature pig as a laboratory animal. *Lab. Anim. Sci.*, 28: 308-313.
- Phillips, R.W., L. Panepinto and D.H. Wild, 1979. Genetic selection for diabetogenic traits in Yucatan miniature swine. *Diabetes*, 28: 1102-1107.
- Phillips, R.W., N. Wesmoreland, L. Panepinto and G.L. Case, 1982. Dietary effects on metabolism of Yucatan Miniature Swine selected for low and high glucose utilization. *J. Nutr.*, 112: 104-111.
- SAS/STAT, 1987. User's Guide for Personal Computers. Version 6th Edn. SAS Institute Inc. Cary, NC.
- Spangler, R., 1980. Characterization of the secretory defect present in glucose intolerant Yucatan miniature swine. Ph.D. Thesis. Colorado State Univ. Fort Collins.
- Steele, N.C., J.P. McMurtry and R.W. Rosebrough, 1985. Endocrine adaptations of periparturient swine to alteration of dietary energy source. *J. Anim. Sci.*, 60: 1260-1268.
- Waltonand, P.E. and T.D. Etherton, 1986. Stimulation of lipogenesis by insulin in swine adipose tissue. Antagonism by porcine growth hormone. *J. Anim. Sci.*, 62: 1584-1595.
- Wangsness, P.J., R.J. Martin and J.H. Gahagan, 1977. Insulin and growth hormone in lean and obese pigs. *Am. J. Physiol. Gastrointest. Liver Physiol.*, 233: G104-G108.
- Wangsness, P.J., W.A. Acker, J.H. Burdette, L.F. Krabill and R. Vasilatos, 1981. Effect of fasting on hormones and metabolites in plasma of fast-growing, lean and slow-growing obese pigs. *J. Anim. Sci.*, 52: 69-74.
- Weiler, U., R. Claus, S. Schnoebelen-Combes and I. Louveau, 1998. Influence of age and genotype on endocrine parameters and growth performance: A comparative study in wild boars, Meishan and Large White boars. *Livest. Prod. Sci.*, 54: 21-31.